

On weak convergence for stochastic evolutionary systems in average principle.

I.V.Samoilenko

Institute of Mathematics,
Ukrainian National Academy of Sciences,
3 Tereshchenkivs'ka, Kyiv, 01601, Ukraine
isamoil@imath.kiev.ua

Abstract: *Weak convergence of the stochastic evolutionary system to the average evolutionary system is proved. The method proposed by R.Liptser in [4] for semimartingales is used. But we apply a solution of singular perturbation problem instead of ergodic theorem.*

Many authors studied the problems of weak convergence for different classes of stochastic processes (see [1-7]). The methods used are also different and depend on the process studied: ergodic theorems, relative compactness, martingales, etc.

We propose to study a stochastic evolutionary system by a combination of two methods. The method proposed by R.Liptser in [4] for semimartingales is combined with a solution of singular perturbation problem instead of ergodic theorem. We use

Theorem [1]. *Let the following conditions hold for a family of Markov processes $\xi^\varepsilon(t), t \geq 0, \varepsilon > 0$:*

C1: *There exists a family of test functions $\varphi^\varepsilon(u, x)$ in $C_0^2(\mathbb{R}^d \times E)$, such that*

$$\lim_{\varepsilon \rightarrow 0} \varphi^\varepsilon(u, x) = \varphi(u),$$

uniformly on u, x .

C2: *The following convergence holds*

$$\lim_{\varepsilon \rightarrow 0} \mathbb{L}^\varepsilon \varphi^\varepsilon(u, x) = \mathbb{L} \varphi(u),$$

uniformly on u, x . The family of functions $\mathbb{L}^\varepsilon \varphi^\varepsilon, \varepsilon > 0$ is uniformly bounded, and $\mathbb{L} \varphi(u)$ and $\mathbb{L}^\varepsilon \varphi^\varepsilon$ belong to $C(\mathbb{R}^d \times E)$.

C3: The quadratic characteristics of the martingales that characterize a coupled Markov process $\xi^\varepsilon(t), x^\varepsilon(t), t \geq 0, \varepsilon > 0$ have the representation $\langle \mu^\varepsilon \rangle_t = \int_0^t \zeta^\varepsilon(s) ds$, where the random functions $\zeta^\varepsilon, \varepsilon > 0$, satisfy the condition

$$\sup_{0 \leq s \leq T} \mathbf{E} |\zeta^\varepsilon(s)| \leq c < +\infty.$$

C4: The convergence of the initial values holds and

$$\sup_{\varepsilon > 0} \mathbf{E} |\zeta^\varepsilon(0)| \leq C < +\infty.$$

Then the weak convergence

$$\xi^\varepsilon(t) \Rightarrow \xi(t), \varepsilon \rightarrow 0,$$

takes place.

The evolutionary system we are going to study is constructed in such a way that the conditions **C1**, **C2** are satisfied as a result of solution of singularly perturbation problem for the system. To verify the conditions **C3**, **C4** we use the method proposed in [4] extending Bogolubov averaging principle for discontinuous semimartingales.

Really, we study a stochastic evolutionary system with the switching ergodic Markov process

$$du_t^\varepsilon/dt = b(u_t^\varepsilon; \mathfrak{x}(t/\varepsilon)), \quad (1)$$

$$u_0^\varepsilon = u.$$

The switching Markov process $\mathfrak{x}(t), t \geq 0$ on the standard (Polish) phase space $(E, \mathcal{E})$ is defined by the generator

$$Q\varphi(x) = \int_E Q(x, dy) [\varphi(y) - \varphi(x)],$$

where

$$Q(x, dy) = q(x)P(x, dy).$$

The stationary distribution $\pi(dx)$ of the ergodic process defines the projector

$$\Pi\varphi(x) = \widehat{\varphi}\mathbf{1}(x), \widehat{\varphi} := \int_E \pi(dx)\varphi(x), \mathbf{1}(x) \equiv 1.$$

We also define a potential R_0 that is, for ergodic Markov process with generator Q and semigroup $P_t, t \geq 0$, a bounded operator

$$R_0 = \int_0^\infty (P_t - \Pi)dt,$$

$$QR_0 = R_0Q = \Pi - I.$$

The velocity function $b(u; x), u \in \mathbb{R}^d, x \in E$ provides a global solution of the deterministic equations

$$du_x(t)/dt = b(u_x(t); x), u_x(0) = u, x \in E.$$

Markov process $u_t^\varepsilon, \mathfrak{x}^\varepsilon(t) := \mathfrak{x}(t/\varepsilon), t \geq 0$ can be characterized by the generator [1, Proposition 3.3]

$$\mathbb{L}^\varepsilon\varphi(u; x) = [\varepsilon^{-1}Q + \mathbb{B}(x)]\varphi(u; x), \quad (2)$$

where the generator $\mathbb{B}(x)\varphi(u) = b(u; x)\varphi'(u)$.

It follows from Proposition 5.6 [1] that the solution of the singular perturbation problem for the generator (2) is given by the relation

$$\mathbb{L}^\varepsilon\varphi^\varepsilon(u; x) = \widehat{\mathbb{B}}\varphi(u) + \varepsilon\theta^\varepsilon(x)\varphi(u) \quad (3)$$

on the functions $\varphi^\varepsilon(u; x) = \varphi(u) + \varepsilon\varphi_1(u; x)$ with $\varphi(u) \in C_0^2(\mathbb{R}^d)$.

Thus, we see from (3) that the solution of singularly perturbation problem for $\mathbb{L}^\varepsilon, \varphi^\varepsilon(u; x)$ satisfies the conditions **C1**, **C2**.

The operator

$$\widehat{\mathbb{B}}\varphi(u) = \widehat{b}(u)\varphi'(u), \widehat{b}(u) = \int_E \pi(dx)b(u; x).$$

The term $\theta^\varepsilon(x) = \mathbb{B}(x)R_0\widehat{\mathbb{B}}(x), \widehat{\mathbb{B}}(x) := \mathbb{B}(x) - \widehat{\mathbb{B}}$ satisfy the condition

$$|\theta^\varepsilon(u)| \leq C < \infty.$$

Remark. The definition of the operator $\widetilde{\mathbb{B}}(x)$ depends on the definition of the potential R_0 . Namely, if $QR_0 = \Pi - I$, as we defined, then $\widetilde{\mathbb{B}}(x) := \mathbb{B}(x) - \widehat{\mathbb{B}}$. If we'd define $QR_0 = I - \Pi$, then we'd have $\widetilde{\mathbb{B}}(x) := \widehat{\mathbb{B}} - \mathbb{B}(x)$.

Let us now regard an average evolutionary system

$$d\widehat{u}_t/dt = \widehat{b}(\widehat{u}_t), \widehat{u}_0 = u. \quad (4)$$

Our aim is to prove a weak convergence of the stochastic evolutionary system (1) to the average evolutionary system (4). To do this, we should show that the conditions **C1-C4** of the Theorem are satisfied. As we showed, the conditions **C1**, **C2** are true due to the solution of singularly perturbation problem for the evolutionary system. Thus, we are going to verify **C3**, **C4**.

Following [4] we demand from the function $b(u; x)$ to satisfy the the following conditions:

I (linear growth)

$$|b(u; x)| \leq L(1 + |u|),$$

II (Lipschitz continuity)

$$|b(u; x) - b(u'; x)| \leq C|u - u'|.$$

To prove the main result we need the following lemma.

Lemma. Under the assumption **I** there exists a constant $k > 0$, independent of ε and dependent on T :

$$\mathbf{E} \sup_{t \leq T} |u_t^\varepsilon|^2 \leq k_T.$$

Corollary. (follows from Chebyshev's inequality) Under the assumption **I** compact containment condition (CCC) holds:

$$\lim_{c \rightarrow \infty} \sup_{\varepsilon \leq \varepsilon_0} \mathbf{P}\{\sup_{t \leq T} |u_t^\varepsilon| > c\} = 0.$$

Proof of Lemma: (follows [4]) Let us rewrite the equation (1) in the form:

$$u_t^\varepsilon = u + \int_0^t b(u_s^\varepsilon; \mathfrak{x}(s/\varepsilon)) ds =: u + A_t^\varepsilon.$$

If we put $u_t^* = \sup_{s \leq t} |u_s|$, then

$$((u_t^\varepsilon)^*)^2 \leq 2[u^2 + ((A_t^\varepsilon)^*)^2]. \quad (6)$$

The condition **I** implies that

$$(A_t^\varepsilon)^* \leq L \int_0^t (1 + (u_s^\varepsilon)^*) ds.$$

The last inequality with (6) and Cochy-Bounikovsky inequality $([\int_0^t \varphi(s) ds]^2 \leq t \int_0^t \varphi^2(s) ds)$ implies, with some constants k_1 and k_2 , independent of ε , that

$$\mathbf{E}((u_t^\varepsilon)^*)^2 \leq k_1 + k_2 \int_0^t \mathbf{E}((u_s^\varepsilon)^*)^2 ds,$$

for $t \leq T$.

By Gronwall-Bellman inequality we obtain

$$\mathbf{E}((u_t^\varepsilon)^*)^2 \leq k_1 \exp(k_2 T).$$

Lemma is proved.

Remark. Another way to prove CCC is proposed in [1, Theorem 8.10] and used by other authors [2,3]. They use the function $\varphi(u) = \sqrt{1 + u^2}$ and prove corollary for $\varphi(u_t^\varepsilon)$ by applying the martingale characterization of the Markov process.

This may be easily done, due to specific properties of $\varphi(u)$.

The main result of our work is the following.

Proposition. Under the conditions **I**, **II**

$$\mathbf{P} - \limsup_{\varepsilon \rightarrow 0} \sup_{t \leq T} |u_t^\varepsilon - \hat{u}_t| = 0. \quad (5)$$

Proof of Proposition: As we mentioned, to prove the Proposition we should verify the conditions **C3**, **C4** of the Theorem.

Remark. Condition **C3** of the Theorem means that the quadratic characteristics of the martingale, corresponding to a coupled Markov

process, is relatively compact. The same result follows from the CCC (see Corollary) by [6].

Thus, the condition **C3** follows from the Corollary.

As soon as $u_0^\varepsilon = \widehat{u}_0 = u$, we see from the Lemma that the condition **C4** follows from **I, II**.

Thus, all the conditions of the Theorem 6.3 [1] are satisfied, so (5) is true.

Proposition is proved.

Remark. To compare the results obtained in this work and in the work [4], we should note that R.Liptser uses the following scheme: he first applies the conditions **I, II** to obtain the result of Lemma 1, and CCC follows from this Lemma. We also apply the conditions **I, II** to obtain Lemma 1 and CCC.

Then, by [4], the necessary weak convergence is obtained by the use of ergodic theorem. But we, on the contrary, use singularly perturbation problem, and so the weak convergence follows from the conditions **C1, C2**.

Similar method is used by A.V.Skorokhod in [7,8]. But his condition of weak convergence looks like existence of a generator, equivalent to our $\widehat{\mathbb{B}}$, that corresponds to a functional of the process. This generator should approximate the functional's surplus.

Another method is used in [2]. Here a well-known conditions of compactness are applied to the family of the processes studied. Then the weak convergence follows from compactness and convergence at a "good" class of test-functions.

In [3,5] similar conditions of compactness are used for a family of processes. Then a submartingale, corresponding to the family of processes is constructed. The weak convergence follows then from the compactness and uniqueness of the solution of martingale problem.

References

1. **Koroliuk, V.S., Limnios, N.** Stochastic Systems in Merging Phase Space. *World Scientific Publishers (2005), 330p.*
2. **Ethier, S.N., Kurtz T.G.** Markov Processes: Characterization and convergence. *J. Wiley & Sons, New York, (1986), 529p.*
3. **Свириденко, М.Н.** Мартингальный подход к получению предельных теорем для марковских процессов. *ВИНТИ №37, (1986), 30с.*
4. **Liptser, R.Sh.** The Bogolubov averaging principle for semimartingales. *Proceedings of the Steclov Institute of Mathematics, No.4, (1994), 12p.*
5. **Stroock, D.W., Varadhan, S.R.S.** Multidimensional Diffusion Processes. *Springer-Verlag, Berlin, (1979), 339p.*
6. **Jacod J., Shiryaev A.N.** Limit Theorems for Stochastic Processes. *Springer-Verlang (1987), Berlin, 325p.*
7. **Скороход, А.В.** Асимптотические методы теории стохастических дифференциальных уравнений. *Наукова думка, Киев, (1987), 328с.*
8. **Skorokhod, A.V., Hoppensteadt, F.C., Salehi, H.** Random Perturbation Methods with Applications in Science and Engineering. *Springer (2002), 488p.*